

Bir Bomba Fabrikası Çalışanlarındaki Total Kolesterol Düzeyleri ile Depresif Durum Arasındaki İlişki

The Relationship Between Depressive Status and Total Cholesterol Levels in a Bomb Plant Workers

Bomba Fabrikası Çalışanlarında Depresyon ve Kolesterol Düzeyleri /
Depression and Cholesterol Levels in Bomb Plant Workers

Hakan Boyunaga¹, Levent Kenar², Hatice Keles³, Mahir Gulec⁴, Omer Oguzturk⁵

¹Department of Biochemistry and Clinical Biochemistry, Kırıkkale University Faculty of Medicine, Kırıkkale, ²Department of Biochemistry and Medical NBC Defense, Gulhane School of Medicine, Ankara, ³Departments of Internal Medicine, Kırıkkale University Faculty of Medicine, Kırıkkale, ⁴Department of Public Health, Gulhane School of Medicine, Ankara, ⁵Departments of Psychiatry Kırıkkale University Faculty of Medicine, Kırıkkale, Turkey.

Özet

Amaç

Düşük kolesterol düzeyleri ile agresif davranışlar, depresyon ve intihar vakaları arasındaki ilişkiyi gösteren pek çok çalışma bulunmaktadır.

Gereç ve Yöntemler

Bu çalışmanın amacı ise, bir bomba fabrikasında çalışan 77 erkek işçi ve 80 ofis çalışanındaki depresif durum ile serum total kolesterol düzeyleri arasında bir korelasyon olup olmadığını ortaya koymaktır. Deneklerin depresif durumları Beck Depresyon Derecelendirme Ölçeği ve Zung Kendi Kendine Derecelendirme Depresyon Ölçeği kullanılarak değerlendirilmiştir.

Bulgular

Ofis çalışanlarına göre bomba fabrikası işçilerinde depresyon skorlamasının oldukça yüksek ve total kolesterol düzeylerinin anlamlı bir şekilde düşük olduğu bulunmuştur. Her iki çeşit personelde kolesterol düzeyleri hem Beck hem de Zung ölçeklerine göre bulunan depresyon skorlaması ile negatif korelasyon göstermekteydi. Orta derecede ve ağır depresyon semptomları gösteren bomba fabrikası işçilerinin total kolesterol düzeyleri ise normal ve hafif depresyon vakalarına göre anlamlı derecede düşük bulundu.

Sonuç

Bu veriler, düşük serum total kolesterol düzeylerinin depresif durumun artması riski ile bir ilişkisinin bulunduğunu göstermektedir.

Anahtar Kelimeler

Depresyon, Kolesterol, Bomba Fabrikası.

Abstract

Aim

There is significant amount of evidence linking low cholesterol levels to aggressive behaviors, depression and suicide.

Material and Methods

The objective of this study was to determine whether a correlation exists between serum total cholesterol levels and depressive state in 77 male workers labouring in a bomb plant and in 80 male office personnel. Depressive statuses of the subjects were determined by using Beck Depression Rating Scale and Zung Self-Rating Depression Scale.

Results

Depression scores were significantly higher and total cholesterol levels were significantly lower in bomb factory workers than that of office staff. In both types of workers, cholesterol levels were negatively correlated with depression scores according to both Beck and Zung scales. Total cholesterol levels of bomb factory workers with moderate and severe depression symptoms were significantly lower when compared with normal and mild depression.

Conclusion

These data indicate that low serum total cholesterol level is associated with increased risk of depressive state.

Keywords

Depression, Cholesterol, Bomb Plant.

DOI: 10.4328/JCAM.10.1.14 Received: 02.09.2009 Accepted: 15.10.2009 Printed: 01.01.2010 J.Clin.Anal.Med. 2010;1.1:21-25

Corresponding author: Assist.Prof. Levent Kenar, MD.PhD. Department of Biochemistry and Medical NBC Defense, Gulhane School of Medicine, 06018, Ankara, TURKEY,

Phone: +90 312 3043329, Fax: +90 312 3046019, E-mail: lkenar@gata.edu.tr

Introduction

Despite the fact that most people are worried about having increased cholesterol levels and the medical establishment continues to provide the suppression of cholesterol levels to abnormal low levels, it is not widely known that there is a significant amount of evidence linking low cholesterol levels to aggressive behaviors and depression [1-6]. However, many contradictory results have also been reported recently [7-15]. Therefore, the association between low or lowered

Material and Methods

This study was performed on 77 male workers (mean age 35 years; range 22-47 years) labouring in the bomb plant department of Kirikkale Bomb Factory and on 80 official male personnel (mean age 36 yr; range 21-46 yr), as the control group, who were all on check-up in Hospital of Kirikkale Faculty of Medicine. It should also be noted that there had been an explosion in the mentioned Factory on 3rd July 1997, and 208 persons within the affected area were injured. These control groups were selected from the official employees who work at a governmental institution other than bomb plant. Thus, this group was not under threat of a bomb factory accidental explosion due to physical proximity. Subjects who had systemic disease or were taking any medication that affected lipid metabolism and mood were not included into the study. Control subjects were both from the same cultural and ethnical background and both social and economical statuses of controls were also similar with those of workers. Prior to the study, the aim of the study was explained to the subjects and

Results

There was no significant difference in age between two groups ($P > 0.05$). There was a strong correlation between Beck and Zung depression scores in all subjects included into the study ($r = 0.693$; $p < 0.001$). Total cholesterol levels and mean depression scores in workers and official personnel are presented in Table 1. Total cholesterol levels of the bomb factory workers were found significantly lower than those of the official personnel (Table 1). Mean scores of depression according to both Beck and Zung depression scales were significantly higher in factory workers when compared to official personnel (Figure 1).

Beck and Zung depression scores negatively correlated with total cholesterol levels in factory workers ($r = -0.373$, $P = 0.001$ and $r = -0.468$, $P = 0.000$, re-

cholesterol and aggressive behaviours and depression remains controversial. The aim of the present study was to determine the level of total cholesterol and possible depression in a group of male workers labouring in a bomb plant, who are considered to be under the risk for the development of some degree of depression, and in a group of male official personnel, all were otherwise known to be good in health, and whether low cholesterol level can be a risk factor for the development of depression in these subjects.

informed consent was obtained from each individual. Beck Depression Rating Scale and Zung Self-Rating Depression Scale, which were modified for our country, were applied to all subjects included into the study [16-18]. Subjects were subgrouped regarding the depression levels based on the Beck's score as follows: Normal 0-8, depression symptoms 9-21, and clinical depression 22 and over. Subjects were also subgrouped based on Zung's score as follows: Normal 0-49, mild depression 50-59, moderate depression 60-69, severe depression 70 and over. Blood samples were collected in the morning after a 12-hour fasting, and the levels of total cholesterol were measured in sera on Olympus AU600 auto analyzer using spectrophotometric technique.

Data obtained were analyzed by using SPSS 9.0 software program and variables were compared with chi-square and two-tailed t-tests. Relationships between variables were tested by linear regression analysis. Statistical significance was assumed for $P < 0.05$. Data were reported as the mean \pm SD.




Figure 1. Depression scores according to Beck (\blacktriangledown) and Zung (\square) depression scales in relation to total cholesterol levels in men labouring in a bomb plant.

Table 1. Total cholesterol levels and depression scores of the bomb factory workers and official personnel.

	Bomb factory workers n = 77 Mean ± SD	Official personnel n = 80 Mean ± SD	P value
Total cholesterol (mg/dl)	196 ± 56	224 ± 73	0.008
Beck- depression score	16 ± 7	10 ± 5	0.000
Zung - depression score	54 ± 11	47 ± 10	0.000

Table 2. Total cholesterol levels of the factory workers and official personnel in each depression level according to Beck and Zung depression scales

	Total cholesterol (mg/dl)			
	Factory workers		Official workers	
	n	Mean ± SD	n	Mean ± SD
Zung-depression level				
Normal	25	222 ± 65	47	237 ± 81
Mild	27	199 ± 35	29	216 ± 53
Moderate	18	174 ± 52	4	127 ± 21
Severe	7	148 ± 48		
Df= 83, F= 6.4, p = 0.001				
Beck-depression level				
Normal	14	256 ± 71	35	242 ± 73
Depression symptoms	48	186 ± 38	45	210 ± 72
Clinical depression	15	172 ± 53		

spectively) and also in official personnel (r = -0.267, p = 0.016 and r = -0.298, p = 0.007, respectively) (Figure 2).

Table 2 shows distribution of the both groups of subjects in subgroups determined by their scores according to both depression scales. We used a 2-way ANOVA with group based on profession (factory workers us official) and depression level based on scores from Zung scale (none, mild, moderate, severe). This had revealed a significant main effect for being workers at bomb plant, (F=6.4, df=83, p < 0.05.) However, depression level had no effect on cholesterol levels. In factory workers, when the subgroups determined by the scores according to Beck depression scale were compared, total cholesterol levels of normal subjects were found to be significantly higher than those of the subjects with depressive symptoms




Figure 2. Depression scores according to Beck (▼) and Zung (□) depression scales in relation to total cholesterol levels in men working as official personnel

(n=48, df=60, t=2,7, p< 0.001) and subjects with clinical depression (n=15, df=27, t=3.1, p<0.001) while latter two subgroups were not different from each other in their cholesterol levels (df=61, t=3.9, p>0.05). When the subgroups determined by the scores according to Zung depression scale were compared, total cholesterol levels in normal subjects were found not to be different from subjects with mild depression (p>0.05), but to be significantly higher from those of the subjects with medium and severe depression (p = 0.019 and p = 0.007, respectively). In the group of official personnel, according to Beck depression scale, total cholesterol levels of the normal subjects (n=35) were significantly higher than the subjects with depressive symptoms (n=45, df=78, t=-2.55, p<0.05). According to Zung depression scale total cholesterol levels in normal subjects were not different from those of the subjects with mild depression (p= 0.392), but higher than those of the subjects with moderate depression (p= 0.01). In the group of official personnel, nobody was determined to be clinically or severely depressive state.

Discussion

In the present study, bomb factory workers were found to be at significantly higher risk for the development of some degree of depression in comparison with official personnel. This is important since even very low levels of depression symptoms can affect the working request and cause hesitation during the work and decrease in working efficiency. Also, in the present study, total cholesterol levels were found to be significantly lower in the factory workers and correlated negatively with depression scores, suggesting that low cholesterol level is a risk factor for development of depression. An association between dietary cholesterol and central serotonergic activity and social behavior was demonstrated in monkeys [19]. It is possible that low cholesterol level changes neuron membrane fluidity, thus, increases presynaptic serotonin reuptake and decreases postsynaptic serotonin activity and triggers depression [20]. It was also shown that serotonin levels are also reduced as well as cholesterol levels in men with depressive symptoms suggesting that cholesterol levels can affect the activity of serotonin, a brain neurotransmitter implicated in the regulation of mood [4,21]. There are significant amount of studies reporting that low cholesterol level had negative effects on psychological

status and had associations with aggressive behaviors and depression [1,3,22-30]. Some investigators examined the correlation between low serum total cholesterol and deaths from suicide and found that those with lower total cholesterol concentration had more risk of committing suicide than those with higher cholesterol levels [26, 31-35]. Although medical establishment continues to provide the suppression of cholesterol levels to abnormal low levels to decrease cardiac morbidity and mortality, increase in suicide risk and number of deaths caused by violence attitudes appeared because of reduction in lipid levels were suggested to lower the success of total mortality reduction [12, 31,36,37]. In contrast to earlier studies, recent large trials of statins did not show an increase in suicide mortality [7,8,38]. But further studies are warranted to conclude the possible relationship between low or lowered serum cholesterol and suicidal behavior.

In conclusion, low serum total cholesterol level seems to be associated with an increased risk of depressive state. Periodical measurements of serum total cholesterol level especially for the groups of subjects who are relatively at higher risk for developing some degree of depression should be performed and a low level be taken into consideration as one of the risk factors.

References

1. Freedman DS, Byers T, Barrett DH, et al. Plasma lipid levels and psychological characteristics in men. *American Journal of Epidemiology*, 1995; 141: 507-517.
2. Golomb BA. Cholesterol and violence: is there a connection? *Ann Int Med* 1998; 28: 478-487.
3. Suarez EC: Relation of trait depression and anxiety to low lipid and lipoprotein concentrations in healthy young adult woman. *Psychosomatic Medicine* 1999; 61: 273-279.
4. Steegmans PH, Hoes AW, Bak AA, Does E, Grobbee DE. Higher Prevalence of Depressive Symptoms in Middle Age Men With low Serum Cholesterol Levels. *Psychosomatic Medicine*, 2000; 62: 205-211.
5. Rafter D. Biochemical markers of anxiety and depression. *Psychiatry Research*, 2001;103: 93-96.
6. Colin A, Reggers J, Castronovo V, Anseau M. Lipids, depression and suicide. *Encephale*, 2003; 29: 49-58.
7. Muldoon MF, Barger SD, Ryan CM, Flory JD, Lehoczyk JP, Matthews KA, Manuck SB. Effects of lovastatin on cognitive function and psychological well-being. *Am J Med*, 2000; 108: 538-546.
8. Stewart RA, Sharples KJ, North FM, Menkes DB, Baker J, Simes J. Long-term assessment of psychological well-being in a randomized placebo-controlled trial of cholesterol reduction with pravastatin. *The LIPID Study Investigators. Archives of Internal Medicine*, 2000; 160: 3144-3152.
9. Manfredini R, Caracciolo S, Salmi R, et al. The association of low serum cholesterol with depression and suicidal behaviours: new hypotheses for the missing link. *J Int Med Res*, 2000;28: 247-257.
10. Coutu MF, Dupuis G, D'Antono B. The impact of cholesterol lowering on patients' mood. *Journal of Behavioral Medicine*, 2001; 24: 517-536.
11. Nakao M, Ando K, Nomura S, et al. Depressive mood accompanies hypercholesterolemia in young Japanese adults. *Japanese Heart Journal*, 2001; 42, 739-748.
12. Jacobs D, Blackburn H, Higgins M, et al. Report on the conference on low blood cholesterol: mortality associations. *Circulation*, 1992; 86: 1046-1060.
13. Ledochowski M, Murr C, Sperner-Unterweger B, Neurauter G, Fuchs D. Association between increased serum cholesterol and signs of depressive mood. *Clin Chem Lab Med*, 2003; 41: 821-824.
14. Ranjekar PK, Hinge A, Hegde MV, et al. Decreased antioxidant enzymes

- and membrane essential polyunsaturated fatty acids in schizophrenic and bipolar mood disorder patients. *Psychiatry Research*, 2003; 121(2):109-22.
15. Deisenhammer EA, Kramer-Reinstadler K, Liensberger D, et al. No evidence for an association between serum cholesterol and the course of depression and suicidality. *Psychiatry Research*, 2004; 121(3): 253-61.
 16. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of General Psychiatry*, 1961; 4: 562-571.
 17. Beck AT. The depression inventory. In: Williams TA, Katz MM, Shield JA, eds. *Anxiety: Current trends in theory and research*. New York: Academic Press, 1972: 343-354.
 18. Zung WWK. A self-rating depression scale. *Archives of General Psychiatry*, 1965;12: 63-70.
 19. Kaplan JR, Shively CA, Fontenot MB, et al. Demonstration of an association among dietary cholesterol, central serotonergic activity, and social behavior in monkeys. *Psychosomatic Medicine*, 1994; 56: 479-484.
 20. Diebold K, Michel G, Schweizer J, et al. Are psychoactive drug induced changes in plasma lipid and lipoprotein levels of significance for clinical remission in psychiatric disorders? *Pharmacopsychiatry*, 1998; 31: 60-67.
 21. Papakostas GI, Petersen T, Mischoulon D, et al. Serum cholesterol and serotonergic function in major depressive disorder. *Psychiatry Research*, 2003; 118(2) : 137-45.
 22. Stewart MA, Stewart SG. Serum cholesterol in antisocial personality. A failure to replicate earlier findings. *Neuropsychobiology*, 1981; 7: 9-11.
 23. Van Doornen LJP, Van Blokland R. Serum cholesterol: sex specific psychological correlates during rest and stress. *Journal of Psychosomatic Research*, 1987; 31: 239-249.
 24. Fowkes FG, Leng GC, Donnan PT, et al. Serum cholesterol, triglycerides and aggression in the general population. *Lancet*, 1992; 340: 995-998.
 25. Lindberg G, Larsson G, Setterlind S, Rastam L: Serum lipids and mood in working men and women in Sweden. *J Epidemiol Commun Health*, 1994; 48:360-363.
 26. Glueck CJ, Kuller FE, Hamer T, et al. Hypocholesterolemia, hypertriglyceridemia, suicide, and suicide ideation in children hospitalized for psychiatric diseases. *Pediatric Research*, 1994; 35: 602-610.
 27. Horsten M, Wamala SP, Vingerhoest A, Orth-Gomer K. Depressive symptoms, social support and lipid profile in healthy middle aged women. *Psychosomatic Medicine*, 1997; 59: 521-528.
 28. Huang TL, Wu SC, Chiang YS, Chen JF. Correlation between serum lipid, lipoprotein concentrations and anxious state, depressive state or major depressive disorder. *Psychiatry Research*, 2003; 118(2) : 147-53.
 29. Wells AS, Reasd NW, Laugharne JD, Ahluvalia NS. Alteration in mood after changing to a low fat diet. *British Journal of Nutrition*, 1998; 79: 23-30.
 30. Pozzi F, Troisi A, Cerilli M, et al. Serum cholesterol and impulsivity in a large sample of healthy young men. *Psychiatry Research*, 2003; 120(3) : 239-45.
 31. Engelberg H. Low serum cholesterol and suicide. *Lancet*, 1992; 339: 727-729.
 32. Ellison LF, Morrison HI. Low Serum Cholesterol Concentration and Risk of Suicide. *Epidemiology*, 2001; 12: 168-172.
 33. Guillem E, Pelissolo A, Notides C, Lepine JP. Relationship between attempted suicide, serum cholesterol level and novelty seeking in psychiatric in-patients. *Psychiatry Research*, 2002; 112(1) : 83-8.
 34. Atmaca M, Kuloglu M, Tezcan E, et al. Serum leptin and cholesterol values in suicide attempters. *Neuropsychobiology*, 2002: 45:124-127.
 35. Lee HJ, Kim YK. Serum lipid levels and suicide attempts. *Acta Psychiatrica Scandinavica*, 2003;108: 215-221.
 36. Wing RR, Matthews KA, Kuller LH, Meilahn EN, Plantiga P. Waist to hip ratio in middle-aged women. Associations with behavioral and psychosocial factors and with changes in cardiovascular risk factors. *Arteriosclerosis Thrombosis*, 1991; 11: 1250-1257.
 37. Smith GD, Shipley MJ, Marmot MG, Rose G. Plasma cholesterol concentrations and mortality. The Whitehall Study. *JAMA*, 1992; 267: 70-76.
 38. Hyyppa MT, Kronholm E, Virtanen A, Leino A, Jula A. Does simvastatin affect mood and steroid hormone levels in hypercholesterolemic men? A randomized double-blind trial. *Psychoneuroendocrinology*, 2003; 28: 181-194.